

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/338176805>

Gentiana urnula Harry Sm. (Gentianaceae), a new record for the flora of Arunachal Pradesh, India

Article in *Journal of Threatened Taxa* · December 2019

DOI: 10.11609/jott.4420.11.15.15083-15086

CITATIONS

0

READS

230

4 authors:



K. S. Kanwal

GB Pant Institute of Himalayan Environment and Development

62 PUBLICATIONS 126 CITATIONS

SEE PROFILE



Umeshkumar L. Tiwari

Botanical Survey of India

62 PUBLICATIONS 135 CITATIONS

SEE PROFILE



Lod Yama

APSCS&T DBT- Centre of Excellence for Bioresources and Sustainable Developme...

7 PUBLICATIONS 9 CITATIONS

SEE PROFILE




Mahendra S. Lodhi

G.B. Pant National Institute of Himalayan Environment

39 PUBLICATIONS 204 CITATIONS

SEE PROFILE



The Journal of Threatened Taxa (JoTT) is dedicated to building evidence for conservation globally by publishing peer-reviewed articles online every month at a reasonably rapid rate at www.threatenedtaxa.org. All articles published in JoTT are registered under [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/) unless otherwise mentioned. JoTT allows unrestricted use, reproduction, and distribution of articles in any medium by providing adequate credit to the author(s) and the source of publication.

Journal of Threatened Taxa

Building evidence for conservation globally

www.threatenedtaxa.org

ISSN 0974-7907 (Online) | ISSN 0974-7893 (Print)

NOTE

***GENTIANA URNULA* HARRY SM. (GENTIANACEAE), A NEW RECORD FOR THE FLORA OF ARUNACHAL PRADESH, INDIA**

Khilendra Singh Kanwal, Umeshkumar Lalchand Tiwari, Lod Yama & Mahendra Singh Lodhi

26 December 2019 | Vol. 11 | No. 15 | Pages: 15083–15086

DOI: 10.11609/jott.4420.11.15.15083-15086



For Focus, Scope, Aims, Policies, and Guidelines visit <https://threatenedtaxa.org/index.php/JoTT/about/editorialPolicies#custom-0>

For Article Submission Guidelines, visit <https://threatenedtaxa.org/index.php/JoTT/about/submissions#onlineSubmissions>

For Policies against Scientific Misconduct, visit <https://threatenedtaxa.org/index.php/JoTT/about/editorialPolicies#custom-2>

For reprints, contact <ravi@threatenedtaxa.org>

The opinions expressed by the authors do not reflect the views of the Journal of Threatened Taxa, Wildlife Information Liaison Development Society, Zoo Outreach Organization, or any of the partners. The journal, the publisher, the host, and the partners are not responsible for the accuracy of the political boundaries shown in the maps by the authors.

Partner



صندوق محمد بن زايد
للمحافظة على
الكائنات الحية

The Mohamed bin Zayed
SPECIES CONSERVATION FUND

Member



Publisher & Host



**GENTIANA URNULA HARRY SM.
(GENTIANACEAE), A NEW RECORD FOR THE
FLORA OF ARUNACHAL PRADESH, INDIA**

**Khilendra Singh Kanwal¹ , Umeshkumar Lalchand
Tiwari² , Lod Yama³  & Mahendra Singh Lodhi⁴ **

^{1,3,4}G.B. Pant National Institute of Himalayan Environment and Sustainable Development, North East Regional Centre, Itanagar, Arunachal Pradesh 791111, India.

²Botanical Survey of India, Arunachal Pradesh Regional Centre, Itanagar, Arunachal Pradesh 791111, India.

¹kskanwal03@gmail.com (corresponding author),

²tigerumesh11@gmail.com, ³lod1437@gmail.com,

⁴mahen29.mail@gmail.com

The genus *Gentiana* L. (Gentianaceae) consists of around 400 species that are mainly distributed in the alpine regions of the world, but also occur in temperate regions of Asia, Europe and the Americas (Ho & Liu 2001; Struwe & Albert 2002; Mabberley 2008; Favre et al. 2016). The Qinghai–Tibet Plateau (QTP) of the Himalaya is considered to be the main centre of diversity for *Gentiana*, hosting around 250 species (Ho & Pringle 1995). The name *Gentiana* is given by Linnaeus after Gentius, the King of Illyria. Gentianas are important medicinal plants in traditional Chinese medicine, and have been used for over 2,000 years for curing various ailments like hypotension, rheumatic pains, fevers and allergic inflammations (Gupta et al. 2012). In India, the genus is mainly distributed in temperate, sub-alpine, and alpine regions of the Himalaya. A total of 73 taxa (66 species, 4 subspecies and 3 varieties) of *Gentiana* are recognised from India, out of which 31 taxa are recorded

from the eastern Himalayan region whereas 27 taxa are confined to the western Himalaya and only five taxa are described from southern India (Sasidharan 2004; Gupta et al. 2012; Maity 2014; Shabir et al. 2017; Maity & Dey 2017; Maity et al. 2018).

A floristic survey was carried out in Tawang District of Arunachal during 2016–17 for the assessment of floral diversity of high altitude areas. During the collection, *Gentiana urnula* Harry Sm. was recorded from Nagula wetland complex area (27.647°N and 91.861°E at an altitude of 4,000m) of Tawang. The Nagula wetland area is very rich in high altitude floral diversity and little explored at present. This species is very rare and endemic to the eastern Himalaya. *Gentiana urnula* is an important medicinal plant and mostly used in Tibetan medicinal system for the treatment of diarrhoea, dysentery, food poisoning and common cold. The identification of the species was confirmed through the consultation of type specimens, the protologue description of the species and consultation of literature (Hooker 1882; Hara 1965, 1975; Polunin & Stainton 1984; Garg 1987; Stainton 1988; Hajara et al. 1996; Ho & Liu 2001; Giri et al. 2008; Chowdhery et al. 2009; Gupta et al. 2012; Maity 2014; Favre et al. 2016; Maity & Dey 2017; Shabir et al. 2017; Shabir et al. 2018; Maity et al. 2018). Furthermore the Herbarium specimens of the Botanical Survey of India (BSI), Itanagar (ARUN) and State Forest Research Institute (SFRI), Itanagar were consulted. International online herbaria and the Global Biodiversity Information Facility



ISSN 0974-7907 (Online)
ISSN 0974-7893 (Print)

**PLATINUM
OPEN ACCESS**



DOI: <https://doi.org/10.11609/jott.4420.11.15.15083-15086>

Editor: K. Ravikumar, Foundation for Revitalisation of Local Health Traditions (FRLHT), Bengaluru, India. **Date of publication:** 26 December 2019 (online & print)

Manuscript details: #4420 | Received 20 July 2018 | Final received 05 November 2019 | Finally accepted 05 December 2019

Citation: Kanwal, K.S., U.L. Tiwari, L. Yama & M.S. Lodhi (2019). *Gentiana urnula* Harry Sm. (Gentianaceae), a new record for the flora of Arunachal Pradesh, India. *Journal of Threatened Taxa* 11(15): 15083–15086. <https://doi.org/10.11609/jott.4420.11.15.15083-15086>

Copyright: © Kanwal et al. 2019. Creative Commons Attribution 4.0 International License. JoTT allows unrestricted use, reproduction, and distribution of this article in any medium by adequate credit to the author(s) and the source of publication.

Funding: Science & Engineering Research Board (SERB), Department of Science & Technology (DST), Government of India.

Competing interests: The authors declare no competing interests.

Acknowledgements: The authors are extremely thankful for the financial support for the project (F.No. EMR/2014/000408) funded by Science & Engineering Research Board (SERB), Department of Science & Technology (DST), Government of India. The authors are thankful to the Director of G.B. Pant National Institute of Himalayan Environment and Sustainable Development and Director of Botanical Survey of India for necessary facilities and encouragement. Special thanks are due to the local communities for sharing valuable information about the Nagula wetland area. The authors are also highly grateful to the Department of Environment and Forest, Government of Arunachal Pradesh, and the Indian Army for giving the necessary permission and support during the field study.



(GBIF) were also explored for species identification and distribution records. *G. urnula* has not been reported earlier from Arunachal Pradesh. The voucher specimen was deposited in the herbarium of the G.B. Pant National Institute of Himalayan Environment and Sustainable Development, (GBP) and Botanical Survey of India (BSI), Itanagar (ARUN) for future references.

Gentiana urnula Harry Smith

Bull. Misc. Inform. Kew. 15: 51. 1961. Ho & Liu, Worldwide Monogr. *Gentiana*: 2001 (Image 1 A, B).

Holotype: Bhutan, Nelli la near Lingshi Dzong, 4,500m, 13 October 1949, Ludlow, Sherriff & Hicks 17458 (BM holotype; EUPS isotypes).

Perennial herbs, sometimes mat forming, 1.5–2.0 cm high. Stems simple or rarely branched, 1 or 2, glabrous. Basal leaves reddish-green, not rosette; cauline leaves crowded upward; petioles 1–1.5 mm long, membranous; lamina truncate-flabelliform, 5–8 mm, truncate or emarginate at apex, abruptly contracted at base, slightly cartilaginous a long margin, papillate near base only; mid-vein cartilaginous and crested, vein 1, upper most pair of leaves often sessile. Flowers terminal, solitary or 2, subsessile. Calyx tubular, obconic; lobes 5, leafy, leathery, orbicular; tube 4–6 mm long, membranous; lobes 3–4 mm long, similar to leaves. Corolla pale bluish-purple to pale yellow with blue streaks, campanulate, 2–3 cm long; lobes broadly ovate, 3–4.5 × 2.5–3.5 mm, apex rounded and cuspidate, entire at margins; plicae broadly ovate to subtruncate, 1–2 mm,

entire at margins or denticulate. Stamens inserted in corolla tube; filaments 5.5–8.5 mm long; anthers 2–3.5 mm long, ellipsoid. Style short; stigma with triangular lobes. Capsules 1.5–1.8 cm; ovoid-ellipsoid; gynophores up to 4cm, slender. Seeds ellipsoid, 2–2.5 mm long, dark brown; seed coats with simple pits.

Flowering and Fruiting: July–October.

Distribution: India (Sikkim, Arunachal Pradesh), Bhutan, Nepal, China (SW Qinghai, E Xizang) (Ho & Liu 2001).

Specimen examined: 1013(GBP), 10.viii.2017, Nagula Lake, Tawang District, Arunachal Pradesh, India, 27.647°N, 91.861°E, 4,000m, coll. Lod Yama & KS. Kanwal (Images 2 & 3).

Habitat and Ecology: The plants were found growing in some isolated pockets in Nagula Lake area of western Arunachal Pradesh in alpine meadows and gravel slope at 4,000m altitude. It is facing threats from livestock mainly from trampling by yaks and horses, unregulated tourism and developmental activities which result in habitat destruction and fragmentation in the area. In future, the species may face further threat from climate change due to very limited population size and restricted distribution in the Himalayan region. Therefore, conservation action should be taken for this rare and endemic species before it becomes extinct in this region. Extensive grazing by yaks along with the consequent human intrusion for plant exhibited more pronounced habitat destruction and made the plant status crucial for immediate management intervention. Proper updated information regarding the species is



Image 1. *Gentiana urnula* Harry Sm.: A—plants in natural habitat | B—flowers. © K.S. Kanwal.




Image 2. Distribution of *Gentiana urnula* Harry Sm. near Nagula wetland, Tawang District.

lacking in India or neighbouring countries, especially with regard to ecological and habitat information, thereby creating huge lacuna in the knowledge base. Qualitative and quantitative inventory of the species is urgently needed for evolving a long term conservation plan of the species. In addition to this, in situ and ex situ conservation measures, awareness through educational programmes, and community participation should also be required for the conservation of *Gentiana urnula* in the region.

References

- Chowdhery, H.J., G.S. Giri, G.D. Pal, A. Pramanik & S.K. Das (2009). *Materials for the Flora of Arunachal Pradesh*. Vol. III. Botanical Survey of India, Calcutta, 357pp.
- Clarke, C.B. (1883). Gentianaceae, pp. 108–109. In: Hooker, J.D. (ed.). *The Flora of British India*. Vol. 4. L. Reeve & Co., London.
- Eflora (2018). http://www.efloras.org/florataxon.aspx?flora_id=2&taxon_id=200018121. Accessed on 13 July 2018.
- Favre, A., I. Michalak, C.H. Chen, J.C. Wang, J.R. Matuszak, S. Sun, H. Yuan, Y.M. Yuan, S. Lena, R. Muellner & Alexandra (2016). Out-of-Tibet: the spatio-temporal evolution of *Gentiana* (Gentianaceae). *Journal of Biogeography* 43: 2–12.
- Garg, S. (1987). *Gentianaceae of the North West Himalaya (A Revision)*. Today & Tomorrow's Printers and Publishers, New Delhi, 122pp.
- Giri, G.S., A. Pramanik & H.J. Chowdhery (2008). *Material for the Flora of Arunachal Pradesh*, Vol. 2 (Asteraceae–Ceratophyllaceae). Botanical Survey of India, Kolkata, 498pp.
- Gupta, S.A., S.K. Mukherjee & M. Mondal (2012). A census of *Gentiana* L. in India, systematics of flowering plants, pp. 53–58. In: Maiti, G. & S.K. Mukherjee (eds.). *Multidisciplinary approaches in Angiosperm Systematics*. University of Kalyani, Kalyani, 448pp.
- Hajra, P.K., D.M. Verma & G.S. Giri (1996). *Materials for the Flora of Arunachal Pradesh*, Vol. I. Botanical Survey of India, Dehra Dun, 544pp.
- Hara, H. (1965). New or noteworthy flowering plants from Eastern Himalaya. *Journal of Japanese Botany* 40: 19–22.
- Hara, H. (1975). Gentianaceae, pp. 90–91. In: Ohashi, H. (ed.). *Flora of Eastern Himalaya - 3*, University of Tokyo.
- Hooker, J.D. (1882). *The Flora of British India*, Vol. III. L. Reeve and Co., London, 482–495pp.
- Ho, T.N. & J.S. Pringle (1995). Gentianaceae, pp. 86–87. In: Wu, Z.Y. & P.H. Raven (eds.). *Flora of China (Gentianaceae through Boraginaceae)*. Beijing: Science Press and St. Louis: Missouri Botanical Garden Press. St. Louis, 140pp.
- Ho, T.N., Liu Shang-wu & Wu Ching-ju (1988). Gentianaceae. *Flora Republicae Popularis Sinicae* 62: 1–411.
- Ho, T.N. & S. Liu (2001). *A Worldwide Monograph of Gentiana*. Science Press. Beijing, China, 663pp.
- Mabberley, D.J. (2008). *The Plant Book - A Portable Dictionary of Plants, their Classification and Uses - 3rd Edition*. Cambridge University Press, Cambridge, 354pp.
- Maity, D. (2014). A new species of *Gentiana* L. (Gentianaceae) from Sikkim Himalaya. *Edinburgh Journal of Botany* 71: 289–296. <https://doi.org/10.1017/S096042861400016X>
- Maity, D. & S.K. Dey (2017). Notes on the occurrence of *Gentiana karelinii* Griseb. (Gentianaceae) in India. *Pleione* 11(2): 525–528.
- Maity, D., S.K. Dey, J. Ghosh & M. Midday (2018). *Gentiana arunii* sp. nov.: Tiny remarkable *Gentiana* from Sikkim Himalaya. *Edinburgh Journal of Botany* 75(1): 117–125. <https://doi.org/10.1017/S0960428617000373>
- Polunin, O. & A. Stainton (1984). *Flowers of the Himalaya*. Oxford University Press, New Delhi, 580pp.
- Sasidharan, N. (2004). *Biodiversity Documentation for Kerala, Part 6 - Flowering Plants*. Kerala Forest Research Institute, Peechi, Kerala, 702pp.
- Shabir, M., P. Agnihotri, D. Husain, J.K. Tiwari & T. Husain (2017). On the current status of the genus *Gentiana* L. (Gentianaceae) in India. *Pleione* 11(1): 16–24.
- Shabir, M., P. Agnihotri, J.K. Tiwari & T. Husain (2018). *Gentiana*




Image 3. Herbarium image of *Gentiana urnula* (GBP) [1013].

aperta (Gentianaceae) - a new record to India from Ladakh Himalaya. *Journal of Threatened Taxa* 10(9): 12286–12289. <https://doi.org/10.11609/jott.4233.10.9.12286-12289>

Smith, H. (1961). Problems relating to the *Gentiana cachemirica* of the Flora of British India. *Kew Bulletin* 15(1): 43–55.

Stainton, A. (1988). *Flowers of the Himalaya: A Supplement*. Oxford University Press, Oxford, New Delhi, 128pp.

Struwe, L. & V.A. Albert (2002). *Gentianaceae: Systematics and Natural History*. Cambridge University Press, Cambridge, 664pp.





PLATINUM
OPEN ACCESS



The Journal of Threatened Taxa (JoTT) is dedicated to building evidence for conservation globally by publishing peer-reviewed articles online every month at a reasonably rapid rate at www.threatenedtaxa.org. All articles published in JoTT are registered under [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/) unless otherwise mentioned. JoTT allows unrestricted use, reproduction, and distribution of articles in any medium by providing adequate credit to the author(s) and the source of publication.

ISSN 0974-7907 (Online) | ISSN 0974-7893 (Print)

December 2019 | Vol. 11 | No. 15 | Pages: 14927–15090

Date of Publication: 26 December 2019 (Online & Print)

DOI: 10.11609/jott.2019.11.15.14927-15090

www.threatenedtaxa.org

Peer Commentary

Observations on the ex situ management of the Sumatran Rhinoceros *Dicerorhinus sumatrensis* (Mammalia: Perissodactyla: Rhinocerotidae): present status and desiderata for conservation

– Francesco Nardelli, Pp. 14927–14941

Communications

Revisiting genetic structure of Wild Buffaloes *Bubalus arnee* Kerr, 1792 (Mammalia: Artiodactyla: Bovidae) in Koshi Tappu Wildlife Reserve, Nepal: an assessment for translocation programs

– Ram C. Kandel, Ram C. Poudel, Amir Sadaula, Prakriti Kandel, Kamal P. Gairhe, Chiranjibi P. Pokheral, Siddhartha B. Bajracharya, Mukesh K. Chalise & Ghan Shyam Solanki, Pp. 14942–14954

A review on status of mammals in Meghalaya, India

– Adrian Wansaindor Lyngdoh, Honnavalli Nagaraj Kumara, P.V. Karunakaran & Santhanakrishnan Babu, Pp. 14955–14970

A comparative analysis of hair morphology of wild and domestic ungulate prey species of Leopard *Panthera pardus fusca* (Mammalia: Carnivora: Felidae) from Goa, India

– Bipin S. Phal Desai, Avelyno H. D'Costa & S.K. Shyama, Pp. 14971–14978

Understanding people's perception and attitudes towards mammalian fauna using qualitative data: a case study in Barail Wildlife Sanctuary, India

– Amir Sohail Choudhury, Rofik Ahmed Barbhuiya & Parthankar Choudhury, Pp. 14979–14988

An assessment of bird communities across Ujjani and its five satellite wetlands in Solapur District of Maharashtra, India

– Shraddha Prabhakar Karikar, Subhash Vitthal Mali, Kulkarni Prasad & Aphale Priti, Pp. 14989–14997

Growth rate of captive Gharials *Gavialis gangeticus* (Gmelin, 1789) (Reptilia: Crocodylia: Gavialidae) in Chitwan National Park, Nepal

– Bed Bahadur Khadka & Ashish Bashyal, Pp. 14998–15003

Amphibian abnormalities and threats in pristine ecosystems in Sri Lanka

– G.K.V.P.T. Silva, W.A.D. Mahaulpatha & Ansem de Silva, Pp. 15004–15014

Diversity and distribution of orchids of Goa, Western Ghats, India

– Jeewan Singh Jalal, Pp. 15015–15042

Short Communications

Efficacy of oxytetracycline and levamisole treatment on the gastrointestinal parasites in captive Lions *Panthera leo*

– Dhareppa Ganager, Gotakanapura Sanjeevamurthy Mamatha, Asoor Muralidhara, Nagappa Lakkundi Jaya & Beechagondahalli Papanna Shivashankar, Pp. 15043–15046

First record in 129 years of the Tamil Treebrown *Lethe drypetis todara* Moore, 1881 (Lepidoptera: Nymphalidae: Satyrinae) from Odisha, India by fruit-baiting

– Anirban Mahata, Sudheer Kumar Jena & Sharat Kumar Palita, Pp. 15047–15052

A review of the leafhopper tribe Agalliini (Hemiptera: Cicadellidae: Megophthalminae) with a revised key to the known Pakistani genera and species

– Hassan Naveed, Kamran Sohail, Waqar Islam & Yalin Zhang, Pp. 15053–15060

The windowpane oyster family Placunidae Rafinesque, 1815 with additional description of *Placuna quadrangula* (Philipsson, 1788) from India

– Rocktim Ramen Das, Vijay Kumar Deepak Samuel, Goutham Sambath, Pandian Krishnan, Purvaja Ramachandran & Ramesh Ramachandran, Pp. 15061–15067

Notes

Recent records of the rare Mountain Tortoiseshell *Aglais rizana* (Moore, 1872) (Lepidoptera: Nymphalidae) in the upper Garhwal, western Himalaya, India, after 100 years

– Arun Pratap Singh & Tribhuwan Singh, Pp. 15068–15071

First report of *Dicranocentroides indica* (Handschin, 1929) (Collembola: Paronellidae) from Odisha, India

– Ashirwad Tripathy, Pp. 15072–15073

Additions to the knowledge of darkling beetles (Coleoptera: Tenebrionidae) from the Indo-Burma Biodiversity Hotspot, Meghalaya, India

– Vishwanath Dattatray Hegde, Pp. 15074–15078

Bhutan Asiabell *Codonopsis bhutanica* Ludlow (Asterales: campanulaceae): a new addition to the Indian flora

– Samiran Panday, Vikas Kumar, Sudhansu Sekhar Dash, Bipin Kumar Sinha & Paramjit Singh, Pp. 15079–15082

***Gentiana urnula* Harry Sm. (Gentianaceae), a new record for the flora of Arunachal Pradesh, India**

– Khilendra Singh Kanwal, Umeshkumar Lalchand Tiwari, Lod Yama & Mahendra Singh Lodhi, Pp. 15083–15086

***Carex phacota*, Spreng. (Cyperaceae): a new record for the central Western Ghats of Karnataka, India**

– E.S.K. Udupa, H.U. Abhijit & K.G. Bhat, Pp. 15087–15088

Book review

Compendium of Traded Indian Medicinal Plants

– Reviewed by A. Rajasekaran, Pp. 15089–15090

Partner



صندوق محمد بن زايد
للمحافظة على
الحيوانات النادرة
The Mohamed bin Zayed
SPECIES CONSERVATION FUND

Member



Publisher & Host

